




Excretion. Oral cavity as excretory organ

***Prof. Zaporozhets
T.Viber +380972420098***



consists of the kidneys, ureters, urinary bladder and urethra

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.




Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



Location of Kidneys


The kidneys which are positioned **retroperitoneally** lie on either side of the vertebral column, high on the posterior wall of the abdominal cavity

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



(a)


Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



(b)

Kidney Structure

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.




Functions of the Kidneys

- **removal of metabolic wastes from the blood and excretion to the outside of the body**
- **regulation of red blood cell production, blood pressure, calcium ion absorption, and the volume, composition, and pH of the blood**


Renal Blood Vessels

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.




Renal Blood Vessels

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.




Structure of a Nephron

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.




Glomerular Capsule

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.




Glomerulus

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.




Nephron and Associated Blood Vessels




Juxtaglomerular Apparatus

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.




Cortical and Juxtamedullary Nephrons

- **cortical nephrons**
 - 80% of nephrons
 - located close to the surface of the kidney
- **juxtamedullary nephrons**
 - regulate water balance
 - located near the renal medulla




Blood Supply of Nephron

- The glomerular capillary receives blood from the afferent arteriole and passes it to the efferent arteriole
- The efferent arteriole gives rise to the peritubular system, which surrounds the tubule
- Capillary loops called **vasa recta** dip down into the medulla



15 Pathway of Blood Flow Through Kidney and Nephron

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



Urine Formation

- **nephrons remove wastes from the blood and regulate water and electrolyte concentrations**
- **urine is the final product of the processes of:**
 - **glomerular filtration**
 - **tubular reabsorption**
 - **tubular secretion**

Urine Formation

17

- **Glomerular Filtration**

- substances move from blood to glomerular capsule


- **Tubular Reabsorption**

- substances move from renal tubules into blood of peritubular capillaries
- glucose, water, urea, proteins, creatine
- amino, lactic, citric, and uric acids
- phosphate, sulfate, calcium, potassium, and sodium ions

- **Tubular Secretion**

- substances move from blood of peritubular capillaries into renal tubules
- drugs and ions


Glomerular Filtration



Glomerular Filtration

19

- The first step in urine formation is filtration of substances out of the glomerular capillaries into the glomerular capsule
- **Glomerular filtrate** passes through the fenestrae of the capillary endothelium



Glomerular Filtrate and Urine Composition

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

TABLE 20.1 Relative Concentrations of Plasma, Glomerular Filtrate, and Urine Components

| Substance | Plasma | Concentrations (mEq/L) | |
|---|--------|----------------------------|-------|
| | | Glomerular Filtrate | Urine |
| Sodium (Na^+) | 142 | 142 | 128 |
| Potassium (K^+) | 5 | 5 | 60 |
| Calcium (Ca^{+2}) | 4 | 4 | 5 |
| Magnesium (Mg^{+2}) | 3 | 3 | 15 |
| Chloride (Cl^-) | 103 | 103 | 134 |
| Bicarbonate (HCO_3^-) | 27 | 27 | 14 |
| Sulfate (SO_4^{-2}) | 1 | 1 | 33 |
| Phosphate (PO_4^{-3}) | 2 | 2 | 40 |
| (mEq/L (milliequivalents per liter) is a commonly used measure of concentration based on how many charges an ion carries. For a substance with a charge of 1, such as Cl^- , a mEq is equal to a millimole.) | | | |
| Substance | Plasma | Concentrations (mg/100 mL) | |
| | | Glomerular Filtrate | Urine |
| Glucose | 100 | 100 | 0 |
| Urea | 26 | 26 | 1,820 |
| Uric acid | 4 | 4 | 53 |
| Creatinine | 1 | 1 | 196 |

Filtration Pressure and Rate

21

Net Filtration Pressure = force favoring filtration – forces opposing filtration
(glomerular capillary hydrostatic pressure) (capsular hydrostatic pressure and glomerular capillary osmotic pressure)


Glomerular Filtration Rate

- directly proportional to the net filtration pressure

Filtration Pressure and Rate

22

- normally the glomerular net filtration pressure is positive causing filtration
- the forces responsible include hydrostatic pressure and osmotic pressure of plasma and the hydrostatic pressure of the fluid in the glomerular capsule




Net Outward Pressure

| | |
|---|----------|
| Outward force, glomerular hydrostatic pressure | = +60 mm |
| Inward force of plasma colloid osmotic pressure | = -32 mm |
| Inward force of capsular hydrostatic pressure | = -18 mm |
| Net outward pressure | = +10 mm |

Amounts of Glomerular Filtrate and Urine

average amounts over a 24 hour period



Control of Filtration Rate


• Primarily three mechanisms are responsible for keeping the GFR constant

• Increased sympathetic impulses decrease GFR by causing afferent arterioles to constrict

• **Renin-angiotensin system**


• Autoregulation

Renin-Angiotensin system




Tubular Reabsorption

- transports substances from the glomerular filtrate into the blood within the peritubular capillary



Sodium and Water Reabsorption

- osmosis reabsorbs water in response to active transport reabsorbing sodium and other solutes in the proximal portion of the renal tubule



Sodium and Water Filtration, Reabsorption, and Excretion


Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

TABLE 20.2 Average Values for Sodium and Water Filtration, Reabsorption, and Excretion

| | Amount Filtered per Day | Amount Reabsorbed per Day (%) | Amount Excreted per Day |
|---------------------|------------------------------------|--|------------------------------------|
| Water (L) | 180 | 178.2 (99%) | 1.8 (1%) |
| Na ⁺ (g) | 630 | 626.8 (99.5%) | 3.2 (0.5%) |

Tubular Secretion


- transports substances from the blood within the peritubular capillary into the renal tubule



Tubular Secretion

29

In distal convoluted tubules, potassium ions or hydrogen ions may be passively secreted in response to active reabsorption of sodium ions




Regulation of Urine Concentration and Volume

30

- the distal convoluted tubule and collecting duct are impermeable to water, so water may be excreted as dilute urine


- if ADH is present, these segments become permeable, and water is reabsorbed by osmosis into the hypertonic medullary interstitial fluid



The Countercurrent Multiplier


31

- helps maintain the NaCl concentration gradient in the medullary interstitial fluid



Countercurrent Mechanism of Vasa Recta

- fluid in ascending limb becomes hypotonic as solute is reabsorbed
- fluid in descending limb becomes hypertonic as it loses water by osmosis



Role of ADH in Regulating Urine Concentration and Volume

33

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

TABLE 20.3

Role of ADH in Regulating Urine Concentration and Volume

1. Concentration of water in the blood decreases.
2. Increase in the osmotic pressure of body fluids stimulates osmoreceptors in the hypothalamus.
3. Hypothalamus signals the posterior pituitary gland to release ADH.
4. Blood carries ADH to the kidneys.
5. ADH causes the distal convoluted tubules and collecting ducts to increase water reabsorption by osmosis.
6. Urine becomes more concentrated, and urine volume decreases.

Functions of Nephron Components

34

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

TABLE 20.4 Functions of Nephron Components

| Part | Function |
|---------------------------------|--|
| Renal Corpuscle | |
| Glomerulus | Filtration of water and dissolved substances from the plasma |
| Glomerular capsule | Receives the glomerular filtrate |
| Renal Tubule | |
| Proximal convoluted tubule | Reabsorption of glucose; amino acids; creatine; lactic, citric, uric, and ascorbic acids; phosphate, sulfate, calcium, potassium, and sodium ions by active transport Reabsorption of proteins by endocytosis Reabsorption of water by osmosis Reabsorption of chloride ions and other negatively charged ions by electrochemical attraction Active secretion of substances such as penicillin, histamine, creatinine, and hydrogen ions |
| Descending limb of nephron loop | Reabsorption of water by osmosis |
| Ascending limb of nephron loop | Reabsorption of sodium, potassium, and chloride ions by active transport |
| Distal convoluted tubule | Reabsorption of sodium ions by active transport Reabsorption of water by osmosis Active secretion of hydrogen ions Secretion of potassium ions both actively and by electrochemical attraction |
| Collecting Duct | Reabsorption of water by osmosis |

(Note: Although the collecting duct is not anatomically part of the nephron, it is functionally connected.)

Urea and Uric Acid Excretion

35

Urea

- by-product of amino acid catabolism
- plasma concentration reflects the amount of protein in diet
- enters renal tubules through glomerular filtration
- contributes to the reabsorption of water from the collecting duct

Uric Acid

- product of nucleic acid metabolism
- enters renal tubules through glomerular filtration
- most reabsorption occurs by active transport
- ~10% secreted and excreted

Urine Composition

- **about 95% water**
- **usually contains urea, uric acid, and creatinine**
- **may contain trace amounts of amino acids and varying amounts of electrolytes**
- **volume varies with fluid intake and environmental factors**

Renal Clearance

- the rate at which a chemical is removed from the plasma
- tests of renal clearance
 - inulin clearance test
 - creatinine clearance test
 - para-aminohippuric acid (PAH) test
- tests of renal clearance used to calculate glomerular filtration rate

Elimination of Urine

- **nephrons**
- **collecting ducts**
- **renal papillae**
- **minor and major calyces**
- **renal pelvis**
- **ureters**
- **urinary bladder**
- **urethra**
- **outside**

Ureters

- 25 cm long
- extend downward posterior to the parietal peritoneum
- parallel to vertebral column
- in pelvic cavity, join urinary bladder
- wall of ureter
 - mucous coat
 - muscular coat
 - fibrous coat


Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



Urinary Bladder


40

- hollow, distensible, muscular organ located within the pelvic cavity, posterior to the symphysis pubis and inferior to the parietal peritoneum




Urinary Bladder

- the internal floor of the bladder includes a triangular area called the **trigone** which has an opening at each of three angles



Cross Section of Urethra


Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



Urethra

43

- tube that conveys urine from the urinary bladder to the outside of the body



Micturition

44

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

TABLE 20.5 Major Events of Micturition

1. Urinary bladder distends as it fills with urine.
2. Stretch receptors in the bladder wall are stimulated, and they signal the micturition center in the sacral spinal cord.
3. Parasympathetic nerve impulses travel to the detrusor muscle, which responds by contracting rhythmically.
4. The need to urinate is urgent.
5. Voluntary contraction of the external urethral sphincter and inhibition of the micturition reflex by impulses from the brainstem and the cerebral cortex prevent urination.
6. Following the decision to urinate, the external urethral sphincter is relaxed, and impulses from the pons and the hypothalamus facilitate the micturition reflex.
7. The detrusor muscle contracts, and urine is expelled through the urethra.
8. Neurons of the micturition reflex center fatigue, the detrusor muscle relaxes, and the bladder begins to fill with urine again.

Life-Span Changes

- **kidneys appear scarred and grainy**
- **kidney cells die**
- **by age 80, kidneys have lost a third of their mass**
- **kidney shrinkage due to loss of glomeruli**
- **proteinuria may develop**
- **renal tubules thicken**
- **harder for kidneys to clear certain substances**
- **bladder, ureters, and urethra lose elasticity**
- **bladder holds less urine**

Clinical Application

46

Glomerulonephritis

- inflammation of glomeruli
- may be acute or chronic
- acute glomerulonephritis usually occurs as an immune reaction to a *Streptococcus* infection
- antigen-antibody complexes deposited in glomeruli and cause inflammation
- most patients recover from acute glomerulonephritis
- chronic glomerulonephritis is a progressive disease and often involves diseases other than that caused by *Streptococcus*
- renal failure may result from chronic glomerulonephritis